

with respect to t_1 , gives a one-dimensional spectrum of full intensity centerband peaks, whatever the value of a .

In its simplest form, our method does not make full use of the available signal energy. Indeed, a spectrum $S(\omega_1, t_2=0)$ is identical to the *integral* over ω_2 of the two-dimensional spectrum $S(\omega_1, \omega_2)$. The integration adds up unwanted noise in addition to the signal energy which is concentrated in the sidebands at (known) equidistant frequencies in ω_2 . More efficient use of the signal energy could in principle be made by sampling all rotary echoes in t_2 and co-adding the signals so obtained.

Experiments were performed on powdered magnesium pyrophosphate, $Mg_2P_2O_7$, which features two isotropic ^{31}P shifts separated by 750 Hz at 121.5 MHz on our Bruker MSL 300 spectrometer. Owing to the high values of the two different ratios a (for $\omega_r/2\pi = 2$ kHz), a conventional TOSS spectrum (Figure 2b) shows severe intensity losses. In contrast, the spectrum acquired using the new method (but with a 90° excitation pulse instead of cross-polarization) has two equal intensity lines, as expected for a method which quantitatively reflects the stoichiometry of the sites (Figure 2c). A similar spectrum could be obtained by the PASS technique, albeit in a more cumbersome manner.¹⁶

Our new experiment is more powerful than TOSS since it not only suppresses the sidebands completely but also transfers the intensity of each sideband family into the corresponding centerband line. Moreover, in contrast to TOSS, this method is applicable to samples with an anisotropic distribution of crystallites, since the t_1 evolution does not depend on their orientation.¹² We use the acronym TIPSy (totally isotropic spectroscopy) for experiments which incorporate this scheme for effecting evolution under the isotropic shifts alone.

Acknowledgment. We thank Dr. Dieter Theimer for experimental assistance and Dr. Christian Jäger for the magnesium pyrophosphate sample. We also acknowledge the Fonds National Suisse de la Recherche Scientifique and the Royal Society of Great Britain (H.G.).

(16) Dixon, W. T.; Schaefer, J.; Sefcik, M. D.; Stejskal, E. O.; McKay, R. A. *J. Magn. Reson.* 1981, 45, 173.

A One-Step Synthesis of the Ciclamycin Trisaccharide

Subharekha Raghavan and Daniel Kahne*

Department of Chemistry, Princeton University
Princeton, New Jersey 08544

Received October 15, 1992

We have been investigating a glycosylation method that involves the use of anomeric phenyl sulfoxides activated with triflic anhydride¹ or catalytic triflic acid.^{2,3} We now report that by using

(1) (a) Kahne, D.; Walker, S.; Cheng, Y.; Van Engen, D. *J. Am. Chem. Soc.* 1989, 111, 6881. (b) Ikemoto, N.; Schreiber, S. L. *J. Am. Chem. Soc.* 1990, 112, 9657. (c) Yang, D.; Kim, S.-H.; Kahne, D. *J. Am. Chem. Soc.* 1991, 113, 4715. (d) Stork, G.; Kim, G. *J. Am. Chem. Soc.* 1992, 114, 1087. (e) Berkowitz, D. B.; Schulte, G. K.; Danishefsky, S. J. *J. Am. Chem. Soc.* 1992, 114, 4518.

(2) A manuscript describing triflic acid activation of anomeric sulfoxides is in preparation. Other triflic acid and triflic anhydride catalyzed glycosylation reactions have been reported: (a) Leroux, J.; Perlin, A. S. *Carbohydr. Res.* 1976, 47, C8-C10. (b) Kishi, Y. *J. Nat. Prod.* 1979, 42, 549. (c) Pavia, A. A.; Rocheville, J.; Ung, S. N. *Carbohydr. Res.* 1980, 79, 79. (d) Lonn, H. *Glycoconjugate J.* 1987, 4, 117. (e) Mootoo, D. R.; Konradsson, P.; Fraser-Reid, B. *J. Am. Chem. Soc.* 1989, 111, 8540. (f) Evans, D. A.; Kaldor, S. W.; Jones, T. K.; Clardy, J.; Stout, T. J. *J. Am. Chem. Soc.* 1990, 112, 7001. (g) Veeneman, G. H.; van Leeuwen, S. H.; van Boom, J. H. *Tetrahedron Lett.* 1990, 31, 1331.





Figure 1. Ciclamycin 0.

Scheme I^a



^a Conditions: Premixed 1 (3.0 equiv), 2 (2.0 equiv), and 3 (1.0 equiv), $Et_2O-CH_2Cl_2$ (1:1), $HC\equiv CCOOCH_3$ (20.0 equiv), TfOH (0.05 equiv), -78 to -70 °C, 45 min, then quenched with saturated $NaHCO_3$ solution.

the sulfoxide method we are able to construct two glycosidic linkages sequentially in a single reaction. This has allowed us to synthesize the ciclamycin 0 trisaccharide stereoselectively from the component monosaccharides in one step.

This synthetic approach grew out of mechanistic studies on the sulfoxide glycosylation reaction. We have found that the rate-limiting step in the reaction is triflation of the sulfoxide; therefore, the reactivity of the glycosyl donor can be influenced by manipulating the substituent in the para position of the phenyl ring (reactivity order: $OMe > H > NO_2$).^{4,5} For perbenzylated glucose sulfoxides, the difference in reactivity is large enough that the *p*-methoxyphenyl sulfoxide can be selectively activated in the presence of an equimolar amount of the corresponding unsubstituted phenyl sulfoxide, as long as only one-half of an equivalent of activating agent is present. Both sulfoxides react, presumably,

(3) O-Glycosylation based on activation of 1-thioglycosides and anomeric sulfones has been studied extensively: (a) Ferrier, R. J.; Hay, R. W.; Vethaviasar, N. *Carbohydr. Res.* 1973, 27, 55. (b) Mukaiyama, T.; Nakatsuka, T.; Shoda, S. *Chem. Lett.* 1979, 487. (c) Van Cleve, J. W. *Carbohydr. Res.* 1979, 70, 161. (d) Hanessian, S.; Baquet, C.; Lehong, N. *Carbohydr. Res.* 1980, 80, C17. (e) Garegg, P. J.; Henrichson, C.; Norberg, T. *Carbohydr. Res.* 1983, 116, 162. (f) Nicolaou, K. C.; Seitz, S. P.; Papahatjis, D. P. *J. Am. Chem. Soc.* 1983, 105, 2430. (g) Lonn, H. *Carbohydr. Res.* 1985, 139, 105, 115. (h) Andersson, F.; Fugedi, P.; Garegg, P. J.; Nashed, M. *Tetrahedron Lett.* 1986, 27, 3919. (i) Sato, S.; Mori, M.; Ito, Y.; Ogawa, T. *Carbohydr. Res.* 1986, 155, C6. (j) Pozsgay, V.; Jennings, H. J. *J. Org. Chem.* 1987, 52, 4653. (k) Murata, S.; Suzuki, T. *Chem. Lett.* 1987, 849. (l) Ito, Y.; Ogawa, T. *Tetrahedron Lett.* 1988, 29, 1061. (m) Brown, D. S.; Ley, S.; Vile, S. *Tetrahedron Lett.* 1988, 4873. (n) Dasgupta, F.; Garegg, P. J. *Carbohydr. Res.* 1988, 177, C13. (o) Kihlberg, J. O.; Leigh, D. A.; Bundle, D. R. *J. Org. Chem.* 1990, 55, 2860.

(4) The mechanistic studies will be described in a full article. S. Raghavan and D. Kahne, manuscript in preparation.

(5) Roy, R.; Andersson, F. O.; Letellier, M. *Tetrahedron Lett.* 1992, 33, 6053.

sequentially, when a full equivalent of activating agent is added. In addition, while silyl ethers are stable to Tf_2O /base activation of anomeric sulfoxides, they are good glycosyl acceptors when catalytic triflic acid is the activating agent; however, they react more slowly than unprotected alcohols because they must be unmasked to couple. The ability to manipulate the reactivity of both the glycosyl donors and the glycosyl acceptors suggested the novel strategy of synthesizing a complicated trisaccharide in one step.

We chose the ciclamycin O trisaccharide as a target for two reasons. First, we are interested in the mode of action of ciclamycin O (Figure 1) and it is difficult to obtain from natural sources.⁶ Second, and more germane to the point here, Danishefsky has published an impressive stepwise synthesis of this trisaccharide using the best available methods for 2-deoxy oligosaccharide synthesis. The Danishefsky synthesis therefore sets a high standard to evaluate the efficacy of our one-step approach.^{7,8}

The ciclamycin O trisaccharide was synthesized stereospecifically from the readily available monosaccharides **1**, **2**, and **3** as shown in Scheme I.⁹⁻¹² The major product, isolated in 25% yield after flash chromatography on silica gel (20% ethyl acetate/petroleum ether), was the desired trisaccharide **5**. No other trisaccharides were produced, and the only other significant coupled product of the reaction was disaccharide **4** (Scheme I, 15% yield), the precursor to the trisaccharide. The yield of trisaccharide **5** in the reaction is limited *not* by any undesired cross-coupling¹³ but by the instability of the glycosyl donors, particularly keto sulfoxide **1** which decomposes readily at room temperature even in the absence of activating agent.

The products of the reaction indicate that glycosylation takes place in a sequential manner as hoped, with *p*-methoxyphenyl sulfoxide **2** activating faster than phenyl sulfoxide **1** and C-4 alcohol **3** reacting faster than C-4 silyl ether **2**. Consistent with this, if the reaction is quenched at -100°C , only the silyl ether of disaccharide **4** can be isolated (60%). We have thus manipulated the reactivity of both the glycosyl donors and the glycosyl acceptors to control the order in which glycosylation takes place.

By way of comparison, the Danishefsky synthesis of the ciclamycin trisaccharide, which makes elegant use of glycal chemistry,¹⁴ requires 14 steps starting from the 3 glycal precursors and

produces the trisaccharide in an overall yield of 9%.⁷ Several steps are needed simply to modulate the reactivity of the donor/acceptor pairs (glycals) to achieve coupling specificity. Although the glycal method remains a very effective method for the construction of 2-deoxy oligosaccharides, the sulfoxide method has allowed us to achieve coupling specificity and construct the ciclamycin trisaccharide stereospecifically in a single step. This has resulted in a dramatic savings in time and labor (less than 3 h from monosaccharides to purified trisaccharide).

Finally, it should be noted that the trisaccharide (**5**) produced in the one-step reaction has an anomeric phenyl sulfide on the A ring. Anomeric phenyl sulfides are stable ("disarmed") to the conditions that activate anomeric phenyl sulfoxides for glycosylation, but they can be readily oxidized under mild conditions.^{5,15} Thus, the sulfoxide glycosylation reaction also lends itself well to an iterative strategy for oligosaccharide synthesis.^{5,14c,d,15,16} The ciclamycin trisaccharide **5** was oxidized to the corresponding sulfoxide in 80% yield (1.2 equiv of *m*-CPBA, CH_2Cl_2 , -78°C , 2 h) and is ready for coupling to the ciclamycin chromophore.

Acknowledgment. This work was supported by the National Institutes of Health.

Supplementary Material Available: MS (FAB) and ^1H and ^{13}C NMR spectra for compound **5** and ^1H NMR spectra for compounds **1-4** (16 pages). Ordering information is given on any current masthead page.

(14) (a) Thiem, J. *Top. Curr. Chem.* **1990**, *154*, 285 and references cited therein. (b) Ito, Y.; Ogawa, T. *Tetrahedron Lett.* **1987**, *28*, 2723. (c) Friesen, R. W.; Danishefsky, S. J. *J. Am. Chem. Soc.* **1989**, *111*, 6656. (d) Halcomb, R. L.; Danishefsky, S. J. *J. Am. Chem. Soc.* **1989**, *111*, 6661. (e) Ramesh, S.; Kalia, N.; Grewal, G.; Franck, R. W. *J. Org. Chem.* **1990**, *55*, 5. (f) Roush, W. R.; Lin, X.-F. *J. Org. Chem.* **1991**, *56*, 5740. (g) Grewal, G.; Kalia, N.; Franck, R. W. *J. Org. Chem.* **1992**, *57*, 2084. (h) Sebesta, D. P.; Roush, W. R. *J. Org. Chem.* **1992**, *57*, 4799.

(15) (a) Mootoo, D. R.; Konradsson, P.; Udodong, U.; Fraser-Reid, B. J. *Am. Chem. Soc.* **1988**, *110*, 5583. (b) Veeneman, G. H.; van Boom, J. H. *Tetrahedron Lett.* **1990**, *31*, 275. (c) Mehta, S.; Pinto, B. M. *Tetrahedron Lett.* **1991**, *32*, 4435.

(16) (a) Nicolaou, K. C.; Dolle, R. E.; Papahatjis, D. P.; Randall, J. L. *J. Am. Chem. Soc.* **1984**, *106*, 4189. (b) Mootoo, D. R.; Konradsson, P.; Fraser-Reid, B. J. *Am. Chem. Soc.* **1989**, *111*, 8540. (c) Barrett, A. G. M.; Bezuidenhout, B. C. B.; Gasielki, A. F.; Howell, A. R.; Russell, M. A. J. *Am. Chem. Soc.* **1989**, *111*, 1392.

(6) Bieber, L. W.; Da Silva Filho, A. A.; De Mello, J. F.; De Lima, O. G.; Do Nascimento, M. S.; Veith, H. J.; Von der Saal, W. *J. Antibiot.* **1987**, *40*, 1335.

(7) Suzuki, K.; Sulikowski, G. A.; Friesen, R. W.; Danishefsky, S. J. *J. Am. Chem. Soc.* **1990**, *112*, 8895.

(8) For other syntheses of trisaccharides related to anthracycline antibiotics, see: (a) El Khadem, H.; Liav, A. *Carbohydr. Res.* **1979**, *74*, 199. (b) Thiem, J.; Gerken, M. *J. Org. Chem.* **1985**, *50*, 954. (c) Monneret, C.; Martin, A.; Pais, M. *J. Carbohydr. Chem.* **1988**, *7*, 417. (d) Thiem, J. *Trends in Synthetic Carbohydrate Chemistry*; Horton, D.; Hawkins, L. D., McGarvey, G. J., Eds.; ACS Symposium Series 386; American Chemical Society: Washington, DC, 1989; Chapter 8. (e) Thiem, J.; Klaffke, W. *J. Org. Chem.* **1989**, *54*, 2006. (f) Kolar, C.; Kneissl, G. *Angew. Chem., Int. Ed. Engl.* **1990**, *29*, 809. (g) Kolar, C.; Kneissl, G.; Wolf, H. *Carbohydr. Res.* **1990**, *208*, 111.

(9) **1** is prepared from L-rhamnose in 60% overall yield. (a) Roth, W.; Pigman, W. *Methods in Carbohydrate Chemistry*; Whistler, R. L., Wolfson, M. L., Eds.; Academic Press Inc.: New York, 1963; Vol. II, p 405. (b) Martin, A.; Pais, M.; Monneret, C. *Carbohydr. Res.* **1983**, *115*, 21. (c) Ferrer, R. J.; Furneaux, R. H. *Carbohydr. Res.* **1976**, *52*, 63. (d) Omura, K.; Swern, D. *Tetrahedron* **1978**, *34*, 1651.

(10) **2** and **3** are prepared from L-fucose with overall yields of 47% and 52%, respectively. (a) Giese, B.; Groninger, K. S.; Witzel, T.; Korth, H.-G.; Sustmann, R. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 233. (b) Reference 9c. (c) Ogawa, T.; Matsui, M. *Carbohydr. Res.* **1978**, *62*, C-1.

(11) The stereoselectivity achieved is a function of the donor-acceptor pairs and the glycosylation conditions (solvent, temperature). We have found that catalytic triflic acid does not anomerize glycosidic linkages at an appreciable rate below -30°C .²

(12) Methyl propiolate is used as a sulfenic acid scavenger. (a) Shelton, J. R.; Davis, K. E. *Int. J. Sulfur Chem.* **1973**, *8*, 205. (b) Block, E. *J. Am. Chem. Soc.* **1972**, *94*, 642.

(13) Less than 5% of the disaccharide from the cross-coupling of phenyl sulfoxide **1** and free alcohol **3** was detected even though **1** is present in excess; no disaccharide from the cross-coupling of **1** and **2** was detected.

Regio- and Stereoselective Formation and Isomerization of 1,3-Cyclohexadienes Catalyzed by Titanium Aryloxy Compounds[†]

Gary J. Balaich and Ian P. Rothwell*

Department of Chemistry, 1393 Brown Building
Purdue University
West Lafayette, Indiana 47907-1393

Received October 6, 1992

One of the conceptually simplest strategies for the synthesis of the 1,3-cyclohexadiene nucleus involves the selective (2 + 2) cycloaddition of 2 equiv of an alkyne with an olefin.¹ The overall formation of three new carbon-carbon bonds during this reaction offers the potential for controlling both the regio- and the stereochemistry of the products.² This communication reports our observations concerning the ability of titanium aryloxy compounds to catalyze this reaction as well as some mechanistic

[†] Dedicated to Professor Bob Squires on the occasion of his 40th birthday.

(1) (a) Shore, N. E. *Chem. Rev.* **1988**, *88*, 1081. (b) Vollhardt, K. P. C. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 539.

(2) (a) Sternberg, E. D.; Vollhardt, K. P. C. *J. Org. Chem.* **1984**, *49*, 1564. (b) Sternberg, E. D.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1980**, *102*, 4839. (c) Wakatsuki, G.; Yamazaki, H. *J. Organomet. Chem.* **1977**, *139*, 169. (d) Gadek, T. R.; Vollhardt, K. P. C. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 802. (e) Wakatsuki, Y.; Kuramitsu, T.; Yamazaki, H. *Tetrahedron Lett.* **1974**, *51/52*, 4549.